

Some Enumerations for Parking Functions

Po-Yi Huang^{a,*}

Jun Ma^{b,†}

Jean Yeh^{c,‡}

^a Department of Mathematics, National Cheng Kung University, Tainan, Taiwan

^b Institute of Mathematics, Academia Sinica, Taipei, Taiwan

^c Department of Mathematics, National Taiwan University, Taipei, Taiwan

Abstract

In this paper, let $\mathcal{P}_{n,n+k;\leq n+k}$ (resp. $\mathcal{P}_{n;\leq s}$) denote the set of parking functions $\alpha = (a_1, \dots, a_n)$ of length n with $n+k$ (respe. n) parking spaces satisfying $1 \leq a_i \leq n+k$ (resp. $1 \leq a_i \leq s$) for all i . Let $p_{n,n+k;\leq n+k} = |\mathcal{P}_{n,n+k;\leq n+k}|$ and $p_{n;\leq s} = |\mathcal{P}_{n;\leq s}|$. Let $\mathcal{P}_{n;\leq s}^l$ denote the set of parking functions $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_{n;\leq s}$ such that $a_1 = l$ and $p_{n;\leq s}^l = |\mathcal{P}_{n;\leq s}^l|$. We derive some formulas and recurrence relations for the sequences $p_{n,n+k;\leq n+k}$, $p_{n;\leq s}$ and $p_{n;\leq s}^l$ and give the generating functions for these sequences. We also study the asymptotic behavior for these sequences.

Keyword: parking function; leading term; asymptotic behavior

*Partially supported by NSC 96-2115-M-006-012

†Email address of the corresponding author: majun@math.sinica.edu.tw

‡jean.yh@ms45.url.com.tw

1 Introduction

Throughout the paper, we let $[n] := \{1, 2, \dots, n\}$ and $[m, n] := \{m, \dots, n\}$. Suppose that n cars have to be parked in m parking spaces which are arranged in a line and numbered 1 to m from left to right. Each car has initial parking preference a_i ; if space a_i is occupied, the car moves to the first unoccupied space to the right. We call $(a_1, \dots, a_n)$ *preference set*. Clearly, the number of preference sets is m^n . If a preference set $(a_1, \dots, a_n)$ satisfies $a_i \leq a_{i+1}$ for $1 \leq i \leq n-1$, then we say that this preference set is *ordered*. If all the cars can find a parking space, then we say the preference set is a *parking function*. If there are exactly k cars which can't be parked, then the preference set is called a *k-flaw preference set*.

Let n , m , s , and k be four nonnegative integers with $1 \leq s \leq m$ and $k \leq n-1$. Suppose there are m parking spaces. We use $\mathcal{P}_{n,m;\leq s;k}$ to denote a set of k -flaw preference sets $(a_1, \dots, a_n)$ of length n satisfying $1 \leq a_i \leq s$ for all i . For $1 \leq l \leq s$, we use $\mathcal{P}_{n,m;\leq s;k}^l$ to denote a set of preference sets $(a_1, \dots, a_n) \in \mathcal{P}_{n,m;\leq s;k}$ such that $a_1 = l$. Let $\mathcal{P}_{n,m;=s;k}$ (resp. $\mathcal{P}_{n,m;=s;k}^l$) be a set of preference sets $(a_1, \dots, a_n) \in \mathcal{P}_{n,m;\leq s;k}$ (resp. $\in \mathcal{P}_{n,m;\leq s;k}^l$) such that $a_j = s$ for some j . Let $p_{n,m;\leq s;k} = |\mathcal{P}_{n,m;\leq s;k}|$, $p_{n,m;\leq s;k}^l = |\mathcal{P}_{n,m;\leq s;k}^l|$, $p_{n,m;=s;k} = |\mathcal{P}_{n,m;=s;k}|$ and $p_{n,m;=s;k}^l = |\mathcal{P}_{n,m;=s;k}^l|$. For any of the above cases, if the parameter k (resp. m) doesn't appear, we understand $k = 0$ (resp. $m = n$); if the parameter m and s are both erased, we understand $s = m = n$.

There are some results about parking functions with $s = m = n$. Riordan introduced parking functions in [11]. He derived that the number of parking functions of length n is $(n+1)^{n-1}$, which coincides with the number of labeled trees on $n+1$ vertices by Cayley's formula. Several bijections between the two sets are known (e.g., see [5, 11, 12]). Furthermore, define a generating function $P(x) = \sum_{n \geq 0} \frac{(n+1)^{n-1}}{n!} x^n$. It is well known that $xP(x)$ is the compositional inverse of the function $\psi(x) = xe^{-x}$, i.e., $\psi(xP(x)) = x$. Riordan concluded that the number of ordered parking functions is $\frac{1}{n+1} \binom{2n}{n}$, which is also equals the number of Dyck path of semilength n . Parking functions have been found in connection to many other combinatorial structures such as acyclic mappings, polytopes, non-crossing partitions, non-nesting partitions, hyperplane arrangements, etc. Refer to [4, 5, 6, 10, 13, 14] for more information.

Any parking function $(a_1, \dots, a_n)$ can be redefined that its increasing rearrangement $(b_1, \dots, b_n)$ satisfies $b_i \leq i$. Pitman and Stanley generalized the notion of parking functions in [10]. Let $\mathbf{x} = (x_1, \dots, x_n)$ be a sequence of positive integers. The sequence $\alpha = (a_1, \dots, a_n)$ is called an $\mathbf{x}$ -parking function if the non-decreasing rearrangement $(b_1, \dots, b_n)$ of α satisfies $b_i \leq x_1 + \dots + x_i$ for any $1 \leq i \leq n$. Thus, the ordinary parking function is the case $\mathbf{x} = (1, \dots, 1)$. By the determinant formula of Gončarov polynomials, Kung and Yan [9] obtained the number of $\mathbf{x}$ -parking functions for an arbitrary $\mathbf{x}$. See also [15, 16, 17] for the explicit formulas and properties for some specified cases of $\mathbf{x}$.

An $\mathbf{x}$ -parking function $(a_1, \dots, a_n)$ is said to be k -leading if $a_1 = k$. Let $q_{n,k}$ denote the number of k -leading ordinary parking functions of length n . Foata and Riordan [5] derived a generating function for $q_{n,k}$ algebraically. Recently, Sen-peng Eu, Tung-shan Fu and Chun-Ju Lai [1] gave a combinatorial approach to the enumeration of $(a, b, \dots, b)$ -parking functions by their leading terms.

Riordan [11] told us the relations between ordered parking functions and Dyck paths. Sen-peng Eu et al. [2, 3] considered the problem of the enumerations of lattice paths with flaws. It is natural to consider the problem of the enumerations of preference sets with flaws. Ordered k -flaw preference sets were studied in [7]. Building on work in this paper, we give enumerations of k -flaw preference sets in [8].

In this paper, we first consider enumerations of parking functions in $\mathcal{P}_{n,m;\leq m}$. When $m \geq n$, Riordan [11] gave a explicit formula $p_{n,m;\leq m} = (m - n + 1)(m + 1)^{n-1}$. We obtain another formula $p_{n,n+k;\leq n+k} = \sum_{r_0+\dots+r_k=n} \binom{n}{r_0,\dots,r_k} \prod_{i=0}^k (r_i + 1)^{r_i-1}$ for any $n \geq 0$ and $k \geq 0$ and find that the sequence $p_{n,n+k;\leq n+k}$ satisfies the recurrence relation $p_{n,n+k;\leq n+k} = \sum_{i=0}^k \binom{n}{i} p_i p_{n-i,n-i+k-1;\leq n-i+k-1}$. When $m < n$, at least $n - m$ cars can't find parking spaces. We conclude that $p_{n,m;\leq m;n-m} = m^n - \sum_{i=0}^{m-2} \binom{n}{i} (i+1)^{i-1} (m-i-1)^{n-i}$ for any $0 \leq m \leq n$.

Then, we focus on the problem of enumerations of parking functions in $\mathcal{P}_{n;\leq s}$. We prove that $p_{n;\leq s} = p_{n,s;\leq s;n-s}$ by a bijection from the sets $\mathcal{P}_{n;\leq s}$ to $\mathcal{P}_{n,s;\leq s;n-s}$ for any $1 \leq s \leq n$. Also we obtain that $p_{n;\leq n-k} = \sum_{i=0}^{k+1} (-1)^i \binom{n}{i} (n-i+1)^{n-i-1} (k+1-i)^i$ for any $0 \leq k \leq n-1$. Furthermore, for any $n \geq k+1$, we derive two recurrence relations $p_{n;\leq n-k} = p_{n;\leq n-k+1} - \sum_{i=1}^k \binom{n}{i} p_{n-i;\leq n-k}$ and

$p_{n;\leq n-k} = (n+1)^{n-1} - \sum_{i=1}^k \binom{n}{i} (k-i+1)(k+1)^{i-1} p_{n-i;\leq n-k}$. Since $p_{n;=n-k+1} = p_{n;\leq n-k+1} - p_{n;\leq n-k}$, we have $p_{n;=n-k} = p_{n;=n-k+1} + \binom{n}{k+1} (n-k)^{n-k-2} - \sum_{i=1}^k \binom{n}{i} p_{n-i;=n-k}$ for any $k \geq 1$ and $n \geq k+1$, with $p_{n;=n} = n^{n-1}$.

Motivated by the work of Foata and Riordan in [5] as well as Sen-Peng Eu et al. in [1], we investigate the problem of the enumerations of some parking functions with leading term l . We derive the formula $p_{n;\leq s}^l = s^{n-1} - \sum_{i=0}^{l-2} \binom{n-1}{i} (s-i-1)^{n-i-1} p_i - \sum_{i=l}^{s-2} \binom{n-1}{i-1} (s-i-1)^{n-i} p_i^l$ for any $1 \leq l \leq s \leq n$. We prove that $p_{n;\leq s}^s = p_{n-1;\leq s}$ by a bijection from the sets $\mathcal{P}_{n;\leq s}^s$ to $\mathcal{P}_{n-1;\leq s}$ for any $1 \leq s \leq n-1$. Furthermore, for any $n \geq k+1$ and $l \leq n-k$, we conclude that $p_{n;\leq n-k}^l = p_{n;\leq n-k+1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;\leq n-k}^l$ and $p_{n;\leq n-k}^l = p_n^l - \sum_{i=1}^k \binom{n-1}{i} (k-i+1)(k+1)^{i-1} p_{n-i;\leq n-k}^l$. Noting that $p_{n;=n-k+1}^l = p_{n;\leq n-k+1}^l - p_{n;\leq n-k}^l$, we obtain $p_{n;=n}^n = p_{n-1}$ and $p_{n;=n-k}^{n-k} = p_{n-1;\leq n-k}$ for any $k \geq 1$. Let $k \geq 1$, then $p_{n;=n-k}^l = p_{n;=n-k+1}^l + \binom{n-1}{k+1} p_{n-k-1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;=n-k}^l$ for any $n \geq k+2$ and $l \leq n-k-1$.

Also we give the generating functions of some sequences. For a fixed $k \geq 0$, we define a generating function $Q_k(x) = \sum_{n \geq 0} \frac{p_{n,n+k;\leq n+k}}{n!} x^n$, then $Q_0(x) = P(x)$, $Q_k(x) = Q_{k-1}(x)P(x)$ for any $k \geq 1$. Let $Q(x, y) = \sum_{k \geq 0} Q_k(x)y^k$, then $Q(x, y) = \frac{P(x)}{1-yP(x)}$.

Let $R_k(x) = \sum_{n \geq k} \frac{p_{n;\leq n-k}}{n!} x^n$ for any $k \geq 0$, then $R_0(x) = P(x)$ and $R_{k+1}(x) = R_k(x) - \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x)$ for any $k \geq 1$, with initial condition $R_1(x) = (1-x)P(x) - 1$. Using this recurrence relation, by induction, we prove that $R_k(x) = P(x) \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i$ for any $k \geq 0$. Let $R(x, y) = \sum_{k \geq 0} R_k(x)y^k$, then $R(x, y) = \frac{P(x)-y}{e^{xy}-y}$.

Let $H_k(x) = \sum_{n \geq k} \frac{p_{n;=n-k}}{n!} x^n$, then $H_k(x) = H_{k-1}(x) + \frac{x^{k+1}}{(k+1)!} P(x) - \sum_{i=1}^k \frac{x^i}{i!} H_{k-i}(x)$ for any $k \geq 2$, with initial conditions $H_0(x) = xP(x) + 1$ and $H_1(x) = P(x)(x - \frac{1}{2}x^2) - x$. In fact, we may prove that for any $k \geq 0$,

$$\begin{aligned} H_k(x) &= P(x) \left[\sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i \right] \\ &\quad - \left[\sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i \right]. \end{aligned}$$

Let $H(x, y) = \sum_{k \geq 0} H_k(x)y^k$, then $H(x, y) = \frac{P(x)(e^{xy}-1)-y^2+y}{y(e^{xy}-y)}$.

Let $L(x) = \sum_{n \geq 1} \frac{p_n^1}{(n-1)!} x^n$ and $T_k(x) = \sum_{n \geq k+1} \frac{p_n^{n-k}}{(n-1)!} x^n$ for any $k \geq 0$, then $L(x)$ and $T_0(x)$ sat-

isfy the differential equations $L(x) + xL'(x) = 2xP(x)$ and $T_0(x) + xT_0'(x) = 2xP(x) + x^2P'(x)$, respectively. Solving these two equations, we have $L(x) = x[P(x)]^2$ and $T_0(x) = xP(x)$. We find that $T_k(x) = T_{k-1}(x) + \frac{(k+1)^{k-1}}{k!}x^{k+1}P(x) - \frac{2(k+1)^{k-2}}{(k-1)!}x^k$ for any $k \geq 1$, equivalently, $T_k(x) = P(x) \sum_{i=0}^k \frac{(i+1)^{i-1}}{i!}x^{i+1} - \sum_{i=1}^k \frac{2(i+1)^{i-2}}{(i-1)!}x^i$. Let $T(x, y) = \sum_{k \geq 0} T_k(x)y^k$, then $T(x, y)$ satisfies the equations $(1-y) \left[T(x, y) + x \frac{\partial T(x, y)}{\partial y} \right] = 2xP(x)P(xy) + x^2P(xy)P'(x) + x^2yP(x)P'(xy) - 2xyP'(xy)$, and $T(x, y) - xP(x) = yT(x, y) + xP(x)[P(xy) - 1] - xy[P(xy)]^2$. The second identity implies $T(x, y) = \frac{xP(xy)[P(x) - yP(xy)]}{1-y}$.

Let $F_{k,s}(x) = \sum_{n \geq s+1} \frac{p_{n; \leq n-k}^{n-s}}{(n-1)!}x^n$ for any $s \geq k \geq 0$, then

$$F_{k,s}(x) = T_s(x) - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!}x^i F_{k-i,s-i}(x)$$

with the initial condition $F_{0,s}(x) = T_s(x)$ and $F_{k,s}(x) = \sum_{i=0}^k \frac{(-1)^i(k+1-i)^i}{i!}x^i T_{s-i}(x)$. Let $F_k(x, y) = \sum_{s \geq k} F_{k,s}(x)y^s$ for any $k \geq 0$, then

$$F_k(x, y) = T(x, y) - \sum_{i=0}^{k-1} T_i(x)y^i - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!}(xy)^i F_{k-i}(x, y)$$

and

$$F_k(x, y) = T(x, y) \sum_{i=0}^k \frac{(-1)^i(k+1-i)^i}{i!}(xy)^i - \sum_{s=0}^{k-1} \sum_{i=0}^{k-1-s} \frac{(-1)^i(k+1-i)^i}{i!}(xy)^i T_s(x)y^s.$$

Let $F(x, y, z) = \sum_{k \geq 0} F_k(x, y)z^k$, then $F(x, y, z) = \frac{T(x, y) - zT(x, yz)}{e^{xyz} - z}$.

Let $D_{k,s}(x) = \sum_{n \geq s+1} \frac{p_{n; = n-k}^{n-s}}{(n-1)!}x^n$ for any $s \geq k \geq 0$, then when $s = k$, we have

$$D_{k,k}(x) = \begin{cases} xp(x) & \text{if } k = 0 \\ x[P(x) - 1] & \text{if } k = 1 \\ P(x) \sum_{i=0}^{k-1} \frac{(-1)^i(k-i)^i}{i!}x^i - \sum_{i=0}^{k-2} \frac{(-1)^i(k-1-i)^i}{i!}x^i & \text{if } k \geq 2 \end{cases}$$

When $s \geq k+1$, $D_{k,s}(x)$ satisfies the following recurrence relation $D_{k,s}(x) = D_{k-1,s}(x) + \frac{x^{k+1}}{(k+1)!}T_{s-k-1}(x) - \sum_{i=1}^k \frac{x^i}{i!}D_{k-i,s-i}(x)$, with initial conditions $D_{0,0}(x) = xP(x)$ and $D_{0,s}(x) =$

$T_{s-1}(x)$ for any $s \geq 1$, equivalently, $D_{k,s}(x) = \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!}T_{s-i}(x) [(k+1-i)^i - (k+2-i)^i]$.

Let $D_k(x, y) = \sum_{s \geq k} D_{k,s}(x)y^s$ for any $k \geq 1$, then $D_k(x, y)$ satisfies the following recurrence

relation

$$\begin{aligned} D_k(x, y) &= D_{k-1}(x, y) - D_{k-1, k-1}(x) y^{k-1} - D_{k-1, k}(x) y^k + D_{k, k}(x) y^k \\ &\quad + \frac{(xy)^{k+1}}{(k+1)!} T(x, y) - \sum_{i=1}^k \frac{(xy)^i}{i!} [D_{k-i}(x, y) - D_{k-i, k-i}(x) y^{k-i}] \end{aligned}$$

with initial condition $D_0(x) = xP(x) + T(x, y)$, and

$$\begin{aligned} D_k(x, y) &= D_{k, k}(x) y^k + \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T(x, y) \\ &\quad - \sum_{s=0}^{k-1} \sum_{i=1}^{k-s} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T_s(x) y^s \end{aligned}$$

. Let $D(x, y, z) = \sum_{k \geq 0} D_k(x, y) z^k$, then $D(x, y, z) = \frac{xe^{xyz}(P(x)-yz)}{e^{xyz}-yz} + \frac{e^{xyz}-1}{z} F(x, y, z)$.

We are interested in the asymptotic behavior for some sequences since there are no explicit formula for some sequences. Let $\mu_k = \lim_{n \rightarrow \infty} \frac{p_{n; \leq n-k}}{(n+1)^{n-1}}$, then $\mu_k = 1 - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} e^{-i} \mu_{k-i}$ for any $k \geq 1$ and $\mu_0 = 1$, equivalently, $\mu_k = \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i}$. Let $\mu(x) = \sum_{k \geq 0} \mu_k x^k$, then $\mu(x) = (e^{\frac{x}{e}} - x)^{-1}$. let $\eta_k = \lim_{n \rightarrow \infty} \frac{p_{n; = n-k}}{(n+1)^{n-1}}$ for any $k \geq 0$ and $\eta(x) = \sum_{k \geq 0} \eta_k x^k$, then $\eta_k = \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i}$ and $\eta(x) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)}$. Let $\tau_l = \lim_{n \rightarrow \infty} \frac{p_n^l}{(n+1)^{n-2}}$ for any $l \geq 1$ and $\tau(x) = \sum_{l \geq 1} \tau_l x^l$, then $\tau_l - \tau_{l+1} = \frac{l^{l-2}}{(l-1)!} e^{-l}$ with the initial condition $\tau_1 = 2$ and $\tau(x) = \frac{\frac{x^2}{e} P(\frac{x}{e}) - 2x}{x-1}$. Let $\rho_{l, k} = \lim_{n \rightarrow \infty} \frac{p_{n; \leq n-k}^l}{(n+1)^{n-2}}$, then $\rho_{l, k} = \rho_{l, k-1} - \sum_{i=1}^k \frac{e^{-i}}{i!} \rho_{l, k-i}$ for any $k \geq 1$, equivalently, $\rho_{l, k} = \rho_{l, 0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i}$. Let $\rho_l(x) = \sum_{k \geq 0} \rho_{l, k} x^k$ and $\rho(x, y) = \sum_{l \geq 1} \rho_l(x) y^l$, then $\rho_l(x) = \rho_{l, 0} (e^{\frac{x}{e}} - x)^{-1}$ and $\rho(x, y) = \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{(y-1)(e^{\frac{x}{e}} - x)}$. Let $\lambda_{l, k} = \lim_{n \rightarrow \infty} \frac{p_{n; = n-k}^l}{(n+1)^{n-2}}$ for any $k \geq 0$ and $l \geq 1$, $\lambda_l(x) = \sum_{k \geq 0} \lambda_{l, k} x^k$ and $\lambda(x, y) = \sum_{l \geq 1} \lambda_l(x) y^l$, then $\lambda_{l, k} = \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \rho_{l, k+1-i}$, $\lambda_l(x) = \frac{e^{\frac{x}{e}} - 1}{x} \rho_l(x)$ and $\lambda(x, y) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)} \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{y-1}$.

We organize this paper as follows. In Section 2, we give some enumerations of parking functions. In Section 3, we obtain the generating functions of some sequences; In Section 4, we study the asymptotic behavior for some sequence in previous sections. In Appendix, we list the values of $P_{n; \leq s}^l$ for $n \leq 7$.

2 Enumerating parking functions

In this section, we give some enumerations of parking functions.

When $m \geq n$, Riordan [11] gave the following explicit formula.

Lemma 2.1. [11] $p_{n,m;\leq m} = (m - n + 1)(m + 1)^{n-1}$ for any $n \leq m$.

Now, for $m \geq n$, we let $k = m - n$. For any $\alpha \in \mathcal{P}_{n,n+k;\leq n+k}$, suppose parking spaces $m_1, \dots, m_k$ are empty in $n + k$ parking spaces. Let

$$r_i = \begin{cases} m_1 - 1 & \text{if } i = 0 \\ m_{i+1} - m_i - 1 & \text{if } i = 1, \dots, k-1 \\ n + k - m_k & \text{if } i = k \end{cases}$$

Clearly, $\sum_{i=0}^k r_i = n$. Let

$$S_i = \begin{cases} \{j \mid 1 \leq a_j \leq m_1 - 1\} & \text{if } i = 0 \\ \{j \mid m_i + 1 \leq a_j \leq m_{i+1} - 1\} & \text{if } i = 1, \dots, k-1 \\ \{j \mid m_k + 1 \leq a_j \leq n + k\} & \text{if } i = k \end{cases}$$

and $\alpha_i = (a_j)_{j \in S_i}$ be a subsequence of α determined by the subscripts in S_i , then $|S_i| = r_i$ and α_i correspond to a parking function of length r_i for each i . There are $\binom{n}{r_0, \dots, r_k}$ ways to choose $r_0, \dots, r_k$ numbers from $[n]$ for the elements of $S_0, \dots, S_k$ respectively. There are $(r_i + 1)^{r_i - 1}$ possibilities for α_i for each $0 \leq i \leq k$. Hence, we obtain the following lemma.

Lemma 2.2. For any $n \geq 0$ and $k \geq 0$, we have

$$p_{n,n+k;\leq n+k} = \sum_{r_0 + \dots + r_k = n} \binom{n}{r_0, \dots, r_k} \prod_{i=0}^k (r_i + 1)^{r_i - 1}.$$

On the other hand, we derive a recurrence relation for $p_{n,n+k;\leq n+k}$.

Lemma 2.3. For $0 \leq k \leq n - 1$, the sequence $p_{n,n+k;\leq n+k}$ satisfies the following recurrence relation

$$p_{n,n+k;\leq n+k} = \sum_{i=0}^n \binom{n}{i} p_i p_{n-i,n+k-i-1;\leq n+k-i-1}.$$

Proof. For any $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_{n,n+k;\leq n+k}$, we suppose the last empty parking space is $n + k - i$. Obviously, $0 \leq i \leq n$. Let $S = \{j \mid n + k - i + 1 \leq a_j \leq n + k\}$ and α_S be a subsequence of α determined by the subscripts in S , then $|S| = i$. Suppose $\alpha_S = (b_1, \dots, b_{i+k})$,

then $(b_1 - n + i, \dots, b_{i+k} - n + i) \in \mathcal{P}_i$. Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T , then $|T| = n - i$ and $\alpha_T \in \mathcal{P}_{n-i, n+k-i-1; \leq n+k-i-1}$.

There are $\binom{n}{i}$ ways to choose i numbers from $[n]$ for the elements in S . There are p_i and $p_{n-i, n+k-i-1; \leq n+k-i-1}$ possibilities for α_S and α_T , respectively. Hence, we have

$$p_{n, n+k; \leq n+k} = \sum_{i=0}^n \binom{n}{i} p_i p_{n-i, n+k-i-1; \leq n+k-i-1}.$$

■

When $n > m$, at least $n - m$ cars can't find parking spaces. We count the number of preference sets in $\mathcal{P}_{n, m; \leq m; n-m}$ for $n \geq m$.

Lemma 2.4. $p_{n, m; \leq m; n-m} = m^n - \sum_{i=0}^{m-2} \binom{n}{i} (i+1)^{i-1} (m-i-1)^{n-i}$ for any $m \leq n$.

Proof. Let A be a set of parking functions $\alpha = (a_1, \dots, a_n)$ of length n such that $a_i \in [m]$ for all i , then $|A| = m^n$. Next, we count the number of the elements α in A such that there are some empty parking spaces. Since there are some empty parking spaces, we may suppose the first parking space which couldn't be occupied is $i+1$. Obviously, $0 \leq i \leq m-2$. Let $S = \{j \mid a_j \leq i, a_j \in \alpha\}$ and α_S be a subsequence of α determined by the subscripts in S , then $|S| = i$ and $\alpha_S \in \mathcal{P}_i$. Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T , then $|T| = n - i$ and $i+2 \leq a_j \leq m$ for any $j \in T$.

There are $\binom{n}{i}$ ways to choose i numbers from $[n]$ for the elements in S . There are $(i+1)^{i-1}$ and $(m-i-1)^{n-i}$ possibilities for the parking function α_S and the preference set α_T , respectively. Hence, we have $p_{n, m; \leq m; n-m} = m^n - \sum_{i=0}^{m-2} \binom{n}{i} (i+1)^{i-1} (m-i-1)^{n-i}$. ■

Now, we consider the problem of enumerations of parking functions in $\mathcal{P}_{n; \leq s}$. Let $p_{0; \leq 0} = 1$ and $p_{n; \leq 0} = 0$ for any $n > 0$.

Lemma 2.5. For any $1 \leq s \leq n$, we have $p_{n; \leq s} = p_{n, s; \leq s; n-s}$.

Proof. For any $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_{n; \leq s}$, if we erase parking spaces $s+1, s+2, \dots, n$, then there are exactly $n-s$ cars which can't be parked since $a_i \leq s$. So, $\alpha \in \mathcal{P}_{n, s; \leq s; n-s}$. Conversely, for any $\alpha \in \mathcal{P}_{n, s; \leq s; n-s}$, since there are s parking spaces, if we add $n-s$ parking

spaces, then all the cars can be parked. We have $\alpha \in \mathcal{P}_{n;\leq s}$. Hence, $p_{n;\leq s} = p_{n,s;\leq s;n-s}$. ■

Using the identity $s^n = \sum_{i=0}^n \binom{n}{i} (i+1)^{i-1} (s-i-1)^{n-i}$ and let $s = n-k$, we obtain the following corollary.

Corollary 2.1. *For any $0 \leq k \leq n-1$, $p_{n;\leq n-k} = \sum_{i=0}^{k+1} (-1)^i \binom{n}{i} (n-i+1)^{n-i-1} (k+1-i)^i$.*

In the following lemma, we give two recurrence relations for $p_{n;\leq n-k}$.

Lemma 2.6. *Let k be an integer with $k \geq 1$. For any $n \geq k$, the sequence $p_{n;\leq n-k}$ satisfies the following two recurrence relations*

$$p_{n;\leq n-k} = p_{n;\leq n-k+1} - \sum_{i=1}^k \binom{n}{i} p_{n-i;\leq n-k} \quad (1)$$

and

$$p_{n;\leq n-k} = (n+1)^{n-1} - \sum_{i=1}^k \binom{n}{i} (k-i+1)(k+1)^{i-1} p_{n-i;\leq n-k}. \quad (2)$$

Proof. First, we consider Identity (1). Given $1 \leq k \leq n$, for any $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_{n;\leq n-k+1}$, let $\lambda(\alpha) = |\{j \in [n] \mid a_j = n-k+1\}|$ and $A_{k,i} = \{\alpha \in \mathcal{P}_{n;\leq n-k+1} \mid \lambda(\alpha) = i\}$. Obviously, $p_{n;\leq n-k} = p_{n;\leq n-k+1} - \sum_{i=1}^k |A_{k,i}|$. It is easy to obtain that $|A_{k,i}| = \binom{n}{i} p_{n-i;\leq n-k}$ for any $i \in [k]$. Hence, we have

$$p_{n;\leq n-k} = p_{n;\leq n-k+1} - \sum_{i=1}^k \binom{n}{i} p_{n-i;\leq n-k}.$$

For deriving Identity (2), we count the number of the parking functions $\alpha = (a_1, \dots, a_n)$ satisfying $a_i > n-k$ for some i . Let $S = \{j \mid a_j \leq n-k, a_j \in \alpha\}$ and α_S be a subsequence of α determined by the subscripts in S . Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T , then $1 \leq |T| \leq k$ and $n-k+1 \leq a_j \leq n$ for any $j \in T$. Suppose $|T| = i$ and $\alpha_T = (b_1, \dots, b_i)$, then $(b_1 - n + k, \dots, b_i - n + k) \in \mathcal{P}_{i,k;\leq k}$.

There are $\binom{n}{i}$ ways to choose i numbers from $[n]$ for the elements in T . Note $\alpha_S \in \mathcal{P}_{n-i,n-k;\leq n-k,k-i}$. There are $p_{n-i,n-k;\leq n-k,k-i}$ and $p_{i,k;\leq k}$ possibilities for α_S and α_T , respectively. Since $p_{i,k;\leq k} = (k-i+1)(k+1)^{i-1}$ for any $i \leq k$, by Lemma 2.5, we have

$$p_{n;\leq n-k} = (n+1)^{n-1} - \sum_{i=1}^k \binom{n}{i} (k-i+1)(k+1)^{i-1} p_{n-i;\leq n-k}.$$

■

Example 2.1. Take $n = 7$ and $k = 4$. By the data in Appendix, we find $p_{7;\leq 3} = 2052$, $p_{3;\leq 3} = 16$, $p_{4;\leq 3} = 61$, $p_{5;\leq 3} = 206$, $p_{6;\leq 3} = 659$. It is easy to check that $p_{7;\leq 3} = 8^6 - \sum_{i=1}^4 \binom{7}{i} (5-i) 5^{i-1} p_{7-i;\leq 3}$.

Corollary 2.2. For any $k \geq 1$ and $n \geq k + 1$, we have

$$p_{n;=n-k} = p_{n;=n-k+1} + \binom{n}{k+1} (n-k)^{n-k-2} - \sum_{i=1}^k \binom{n}{i} p_{n-i;=n-k}$$

with $p_{n;=n} = n^{n-1}$.

Proof. It is easy to check that $p_{n;=n} = \binom{n}{1} p_{n-1} = n^{n-1}$ for any $n \geq 1$. Note that $p_{n;=n-k+1} = p_{n;\leq n-k+1} - p_{n;\leq n-k}$ for any $k \geq 1$. By Lemma 2.6, we obtain the results as desired. ■

In next subsection, we investigate the problem of enumerations of parking functions by leading term. Sen-peng Eu et. al.[1] have deduced the following lemma.

Lemma 2.7. [1] The sequence p_n^l satisfies the following recurrence relations

$$p_n^l - p_n^{l+1} = \binom{n-1}{l-1} l^{l-2} (n-l+1)^{n-l-1} \quad (3)$$

for $1 \leq l \leq n-1$, with the initial condition

$$p_n^1 = 2(n+1)^{n-2}.$$

We consider enumerations of parking functions by leading term in $\mathcal{P}_{n;\leq s}$.

Theorem 2.1. For any $1 \leq l \leq s \leq n$,

$$p_{n;\leq s}^l = s^{n-1} - \sum_{i=0}^{l-2} \binom{n-1}{i} p_i (s-i-1)^{n-i-1} - \sum_{i=l}^{s-2} \binom{n-1}{i-1} p_i^l (s-i-1)^{n-i}.$$

Proof. Let A be the set of the sequences $(l, a_2, \dots, a_n)$ of length n with leading term l such that $a_i \leq s$ for all i , then $|A| = s^{n-1}$. Next, we count the number of the elements α in A which aren't parking functions. Since α isn't parking function, we may suppose the first parking space which couldn't be occupied is $i+1$. Obviously, $0 \leq i \leq s-2$. Let $S = \{j \mid a_j \leq i, a_j \in \alpha\}$ and α_S be a subsequence of α determined by the subscripts in S ,

then $|S| = i$ and $\alpha_S \in \mathcal{P}_i$. Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T , then $|T| = n - i$ and $i + 2 \leq a_j \leq k$ for any $j \in T$. Now, we discuss the following two cases.

Case I. $0 \leq i \leq l - 2$

Obviously, $1 \in T$. There are $\binom{n-1}{i}$ ways to choose i numbers from $[2, n]$ for the elements in S . There are p_i and $(s - i - 1)^{n-i-1}$ possibilities for α_S and α_T , respectively.

Case II. $l \leq i \leq s - 2$

Note $1 \in S$, hence, there are $\binom{n-1}{i-1}$ ways to choose $i - 1$ numbers from $[2, n]$ for the other elements in S . Since the leading term of α_S is m , there are p_i^l and $(s - i - 1)^{n-i}$ possibilities for α_S and α_T , respectively.

So, the number of the elements α in A which aren't parking functions is equal to

$$\sum_{i=0}^{l-2} \binom{n-1}{i} p_i (s - i - 1)^{n-i-1} + \sum_{i=l}^{s-2} \binom{n-1}{i-1} p_i^l (s - i - 1)^{n-i}.$$

Hence,

$$p_{n;\leq s}^l = s^{n-1} - \sum_{i=0}^{l-2} \binom{n-1}{i} p_i (s - i - 1)^{n-i-1} - \sum_{i=l}^{s-2} \binom{n-1}{i-1} p_i^l (s - i - 1)^{n-i}.$$

■

Example 2.2. Take $n = 7$, $s = 6$ and $l = 3$. By the data in Appendix, we find $p_{7;\leq 6}^3 = 23667$, $p_0 = p_1 = 1$, $p_3^3 = 3$, $p_4^3 = 25$. It is easy to check that $p_{7;\leq 6}^3 = 6^6 - \sum_{i=0}^1 \binom{6}{i} p_i (5 - i)^{6-i} - \sum_{i=3}^4 \binom{6}{i-1} p_i^3 (5 - i)^{7-i}$.

Corollary 2.3. $p_{n;\leq n}^n = p_{n-1}$ for any $n \geq 1$. For any $1 \leq s \leq n - 1$, we have $p_{n;\leq s}^s = p_{n-1;\leq s}$.

Proof. Taking $l = s$ in Theorem 2.1, by Lemmas 2.4 and 2.5, we immediately obtain $p_{n;\leq s}^s = p_{n-1;\leq s}$. Also we give this identity a bijection proof. For any $\alpha = (s, a_1, \dots, a_{n-1}) \in \mathcal{P}_{n;\leq s}^s$, it is easy to obtain that $(a_1, \dots, a_{n-1}) \in \mathcal{P}_{n-1;\leq s}$. Obviously, this is a bijection. Hence, $p_{n;\leq s}^s = p_{n-1;\leq s}$ for any $1 \leq s \leq n - 1$. Using the same method, it is easy to prove that $p_{n;\leq n}^n = p_{n-1}$ for any $n \geq 1$. ■

Similar to Lemma 2.5, we have the following results.

Lemma 2.8. $p_{n;\leq s}^l = p_{n,s;\leq s;n-s}^l$ for any $1 \leq s \leq n$.

In the following lemma, we will derive two recurrence relations for $p_{n;\leq n-k}^l$.

Lemma 2.9. Let k be an integer with $k \geq 1$. For any $n \geq k+1$ and $1 \leq l \leq n-k$, the sequence $p_{n;\leq n-k}^l$ satisfies the following recurrence relation

$$p_{n;\leq n-k}^l = p_{n;\leq n-k+1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;\leq n-k}^l \quad (4)$$

and

$$p_{n;\leq n-k}^l = p_n^l - \sum_{i=1}^k \binom{n-1}{i} (k-i+1)(k+1)^{i-1} p_{n-i;\leq n-k}^l. \quad (5)$$

Proof. First, we consider Identity (4). Given $1 \leq k \leq n$, for any $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_{n;\leq n-k+1}^l$, let $\lambda(\alpha) = |\{j \mid a_j = n-k+1\}|$ and $A_{k,i} = \{\alpha \in \mathcal{P}_{n;\leq n-k+1}^l \mid \lambda(\alpha) = i\}$. Obviously, $p_{n;\leq n-k}^l = p_{n;\leq n-k+1}^l - \sum_{i=1}^k |A_{k,i}|$. It is easy to obtain that $|A_{k,i}| = \binom{n-1}{i} p_{n-i;\leq n-k}^l$ for any $i \in [k]$. Hence, we have

$$p_{n;\leq n-k}^l = p_{n;\leq n-k+1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;\leq n-k}^l.$$

For deriving Identity (5), we count the number of the parking functions $\alpha = (a_1, \dots, a_n) \in \mathcal{P}_n^l$ satisfying $a_i > n-k$ for some i . Let $S = \{j \mid a_j \leq n-k, a_j \in \alpha\}$ and α_S be a subsequence of α determined by the subscripts in S . Let $T = [n] \setminus S$ and α_T be a subsequence of α determined by the subscripts in T , then $1 \leq |T| \leq k$ and $n-k+1 \leq a_j \leq n$ for any $j \in T$. Suppose $|T| = i$ and $\alpha_T = (b_1, \dots, b_i)$, then $(b_1 - n + k, \dots, b_i - n + k) \in \mathcal{P}_{i,k;\leq k}$.

Since $1 \in S$, there are $\binom{n-1}{i}$ ways to choose i numbers from $[2, n]$ for the elements in T . Note that $\alpha_S \in \mathcal{P}_{n-i,n-k;\leq n-k,k-i}^l$. There are $p_{n-i,n-k;\leq n-k,k-i}^l$ and $p_{i,k;\leq k}$ possibilities for parking functions α_S and α_T , respectively. By Lemma 2.8, we have

$$p_{n;\leq n-k}^l = p_n^l - \sum_{i=1}^k \binom{n-1}{i} p_{i,k;\leq k} p_{n-i;\leq n-k}^l.$$

■

Example 2.3. Take $n = 7$, $k = 1$ and $l = 3$. By the data in Appendix, we find $p_{7;\leq 6}^3 = 23667$, $p_7^3 = 40953$, $p_{6;\leq 6}^3 = 2881$. It is easy to check that $p_{7;\leq 6}^3 = p_7^3 - \binom{6}{1} p_{6;\leq 6}^3$.

Corollary 2.4. $p_{n;=n}^n = p_{n-1}$ and $p_{n;=n-k}^{n-k} = p_{n-1;\leq n-k}$ for any $k \geq 1$. Let $k \geq 1$, then

$$p_{n;=n-k}^l = p_{n;=n-k+1}^l + \binom{n-1}{k+1} p_{n-k-1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;=n-k}^l$$

for any $n \geq k+2$ and $l \leq n-k-1$.

Proof. Obviously, $p_{n;=n-k}^{n-k} = p_{n;\leq n-k}^{n-k}$. Hence, we easily obtain that $p_{n;=n}^n = p_{n-1}$ and $p_{n;=n-k}^{n-k} = p_{n-1;\leq n-k}$ for any $k \geq 1$. Note that $p_{n;=n-k+1}^l = p_{n;\leq n-k+1}^l - p_{n;\leq n-k}^l$ for $l \leq n-k$. By Lemma 2.9, we obtain $p_{n;=n-k}^l = p_{n;=n-k+1}^l + \binom{n-1}{k+1} p_{n-k-1}^l - \sum_{i=1}^k \binom{n-1}{i} p_{n-i;=n-k}^l$ for any $n \geq k+2$ and $l \leq n-k-1$. ■

3 Generating functions of parking functions

In this section, we give the generating function for some sequences in previous sections.

For a fixed $k \geq 0$, we define a generating functions $Q_k(x) = \sum_{n \geq 0} \frac{p_{n,n+k;\leq n+k}}{n!} x^n$. Let $Q(x, y) = \sum_{k \geq 0} \sum_{n \geq 0} \frac{p_{n,n+k;\leq n+k}}{n!} x^n y^k$.

Theorem 3.1. For any $k \geq 0$, let $Q_k(x)$ be the generating function for $p_{n,n+k;\leq n+k}$. Then $Q_0(x) = P(x)$ and $Q_k(x)$ satisfies the following recurrence relation

$$Q_k(x) = P(x)Q_{k-1}(x).$$

Furthermore, we have

$$Q_k(x) = [P(x)]^{k+1}.$$

Let $Q(x, y) = \sum_{k \geq 0} Q_k(x) y^k$, then

$$Q(x, y) = \frac{P(x)}{1 - yP(x)}.$$

Proof. Clearly, $Q_0(x) = P(x)$. By Lemma 2.3, we have

$$Q_k(x) = \sum_{n \geq 0} \frac{p_{n,n+k;\leq n+k}}{n!} x^n = \sum_{n \geq 0} \sum_{i=0}^n \frac{p_i}{i!} \frac{p_{n-i,n+k-i-1;\leq n+k-i-1}}{(n-i)!} x^n = P(x)Q_{k-1}(x).$$

Therefore, $Q_k(x) = [P(x)]^{k+1}$ for any $k \geq 0$ and $Q(x, y) = \frac{P(x)}{1-yP(x)}$. ■

Now, for a fixed k , we define a generating function $R_k(x) = \sum_{n \geq k} \frac{p_{n;\leq n-k}}{n!} x^n$.

Theorem 3.2. Suppose k is a nonnegative integer. Let $R_k(x)$ be the generating function for $p_{n;\leq n-k}$. Then $R_0(x) = P(x)$ and $R_k(x)$ satisfies the following recurrence relation

$$R_{k+1}(x) = R_k(x) - \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x)$$

for any $k \geq 1$, with initial condition $R_1(x) = (1-x)P(x) - 1$. Furthermore, we have

$$R_k(x) = P(x) \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i. \quad (6)$$

Proof. Clearly, $R_0(x) = P(x)$. Taking $k = 1$ in Identity (1), we obtain that $p_{n;\leq n-1} = p_n - \binom{n}{1} p_{n-1}$. This implies that $R_1(x) = (1-x)P(x) - 1$. For any $k \geq 2$, using Identity (1), we have

$$\begin{aligned} \sum_{n \geq k+1} \frac{p_{n;\leq n-k-1}}{n!} x^n &= \sum_{n \geq k+1} \frac{p_{n;\leq n-k}}{n!} x^n - \sum_{i=1}^{k+1} \sum_{n \geq k+1} \binom{n}{i} \frac{p_{n-i;\leq n-k-1}}{n!} x^n \\ &= \sum_{n \geq k+1} \frac{p_{n;\leq n-k}}{n!} x^n - \sum_{i=1}^{k+1} \frac{x^i}{i!} \sum_{n \geq k+1} \frac{p_{n-i;\leq n-k-1}}{(n-i)!} x^n. \end{aligned}$$

Since $p_{k;\leq 0} = 0$ for any $k \geq 1$, we have $R_{k+1}(x) = R_k(x) - \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x)$.

It is easy to check that Identities (6) hold when $k = 0, 1$. We assume Identities (6) hold for any $k' \leq k$. Then

$$\begin{aligned} R_{k+1}(x) &= R_k(x) - \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x) \\ &= P(x) \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i \\ &\quad - \sum_{i=1}^{k+1} \frac{x^i}{i!} \left[P(x) \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} x^j - \sum_{j=0}^{k-i} \frac{(-1)^j (k+1-i-j)^j}{j!} x^j \right]. \end{aligned}$$

Note that

$$\begin{aligned} &\sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=1}^{k+1} \frac{x^i}{i!} \sum_{j=0}^{k-i} \frac{(-1)^j (k+1-i-j)^j}{j!} x^j \\ &= \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=1}^k \frac{x^i}{i!} \sum_{j=1}^i \binom{i}{j} (i-k-1)^{i-j} \\ &= \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=1}^k \frac{x^i}{i!} [(i-k)^i - (i-k-1)^i] \\ &= \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i. \end{aligned}$$

Hence,

$$R_{k+1}(x) = P(x) \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i - \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i.$$

■

Corollary 3.1. *Let $R(x, y) = \sum_{k \geq 0} R_k(x) y^k$, then*

$$R(x, y) = \frac{P(x) - y}{e^{xy} - y}$$

Proof. By Theorem 3.2, we have $\sum_{k \geq 1} R_{k+1}(x) y^k = \sum_{k \geq 1} R_k(x) y^k - \sum_{k \geq 1} \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x) y^k$. So, we obtain

$$\frac{1}{y} [R(x, y) - R_0(x) - R_1(x)y] = R(x, y) - R_0(x) - \frac{e^{xy} - 1}{y} R(x, y) + x R_0(x).$$

Hence,

$$R(x, y) = \frac{P(x) - y}{e^{xy} - y}.$$

■

For a fixed k , we define a generating function $H_k(x) = \sum_{n \geq k} \frac{p_{n;=n-k}}{n!} x^n$.

Corollary 3.2. *Let $k \geq 0$ and $H_k(x)$ be the generating function for $p_{n;=n-k}$. Then $H_k(x)$ satisfies the following recurrence relation*

$$H_k(x) = H_{k-1}(x) + \frac{x^{k+1}}{(k+1)!} P(x) - \sum_{i=1}^k \frac{x^i}{i!} H_{k-i}(x),$$

for any $k \geq 2$, with initial conditions $H_0(x) = xP(x) + 1$ and $H_1(x) = P(x)(x - \frac{1}{2}x^2) - x$, equivalently,

$$\begin{aligned} H_k(x) &= P(x) \left[\sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i \right] \\ &\quad - \left[\sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i \right]. \end{aligned}$$

Proof. By Corollary 2.2, for any $k \geq 1$, we have

$$H_k(x) = H_{k-1}(x) - \frac{p_{k-1;=0}}{(k-1)!} x^{k-1} + \frac{x^{k+1}}{(k+1)!} P(x) - \sum_{i=1}^k \frac{x^i}{i!} H_{k-i}(x)$$

and $H_0(x) = xP(x) + 1$. Taking $k = 1$, we obtain $H_1(x) = P(x)(x - \frac{1}{2}x^2) - x$. Since $p_{k-1;=0} = 0$ for any $k \geq 2$, we have

$$H_k(x) = H_{k-1}(x) + \frac{x^{k+1}}{(k+1)!}P(x) - \sum_{i=1}^k \frac{x^i}{i!}H_{k-i}(x).$$

By Lemma 2.6, since $p_{n;=n-k+1} = p_{n;\leq n-k+1} - p_{n;\leq n-k}$ for any $k \geq 1$, we have

$$p_{n;=n-k+1} = \sum_{i=1}^k \binom{n}{i} p_{n-i;\leq n-k}.$$

So, $H_{k-1}(x) - \frac{p_{k-1;=0}}{(k-1)!}x^{k-1} = \sum_{i=1}^k \frac{x^i}{i!}R_{k-i}(x)$. Since $p_{k;=0} = 0$ for any $k \geq 1$, we have

$$\begin{aligned} H_k(x) &= \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x) \\ &= \sum_{i=1}^{k+1} \frac{x^i}{i!} \left[P(x) \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} x^j - \sum_{j=0}^{k-i} \frac{(-1)^j (k+1-i-j)^j}{j!} x^j \right] \\ &= P(x) \left[\sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i \right] \\ &\quad - \left[\sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i \right] \end{aligned}$$

■

Corollary 3.3. Let $H(x, y) = \sum_{k \geq 0} H_k(x) y^k$, then

$$H(x, y) = \frac{P(x)(e^{xy} - 1) - y^2 + y}{y(e^{xy} - y)}.$$

Proof. Note that $H_0(x) = 1 + xP(x)$ and $H_k(x) = \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x)$ for any $k \geq 1$. Hence,

$$\begin{aligned} H(x, y) &= H_0(x) + \sum_{k \geq 1} \sum_{i=1}^{k+1} \frac{x^i}{i!} R_{k+1-i}(x) y^k \\ &= 1 + \frac{e^{xy} - 1}{y} R(x, y) \\ &= \frac{P(x)(e^{xy} - 1) - y^2 + y}{y(e^{xy} - y)}. \end{aligned}$$

■

Let $\psi(x) = xe^{-x}$, it is well known that $xP(x)$ is the inverse function of $\psi(x)$, i.e., $\psi(xP(x)) = x$. Hence, we have $\psi'(xP(x))[P(x) + xP'(x)] = 1$, this implies that $P(x)$ satisfies the differential equation $P'(x) = [P(x)]^2 + xP(x)P'(x)$.

Let $L(x) = \sum_{n \geq 1} \frac{p_n^1}{(n-1)!} x^n$. For a fixed $k \geq 0$, we define a generating function $T_k(x) = \sum_{n \geq k+1} \frac{p_n^{n-k}}{(n-1)!} x^n$ and $T(x, y) = \sum_{k \geq 0} T_k(x) y^k$.

Lemma 3.1. *Let $T_0(x)$ and $L(x)$ be the generating functions for the sequences p_n^n and p_n^1 , respectively, then $L(x)$ and $T_0(x)$ satisfy the following differential equations*

$$L(x) + xL'(x) = 2xP'(x)$$

and

$$T_0(x) + xT_0'(x) = 2xP(x) + x^2P'(x),$$

respectively. Furthermore, we have $L(x) = x[P(x)]^2$ and $T_0(x) = xP(x)$.

Proof. Lemma 2.7 tells us that $p_n^1 = 2(n+1)^{n-2}$. By comparing coefficient, it is easy to prove that $L(x)$ satisfies the differential equation $L(x) + xL'(x) = 2xP'(x)$. Since $P'(x) = [P(x)]^2 + xP(x)P'(x)$ and $L(0) = 0$, solving this equation, we have $L(x) = x[P(x)]^2$.

In Lemma 2.7, take $l = n - k$, then

$$p_n^{n-k} - p_n^{n-k+1} = \binom{n-1}{k} (n-k)^{n-k-2} (k+1)^{k-1} \quad (7)$$

for $1 \leq k \leq n-1$. So, we have

$$p_n^1 - p_n^n = \sum_{k=1}^{n-1} \binom{n-1}{k} (n-k)^{n-k-2} (k+1)^{k-1}.$$

Therefore, $\sum_{n \geq 1} \frac{p_n^1}{(n-1)!} x^n - \sum_{n \geq 1} \frac{p_n^n}{(n-1)!} x^n = \sum_{n \geq 1} \sum_{k=1}^{n-1} \frac{(n-k)^{n-k-2} (k+1)^{k-1}}{(n-k-1)! k!} x^n$. We obtain that $T_0(x) = L(x) + xP(x) - x[P(x)]^2$. Since $L(x) + xL'(x) = 2xP'(x)$ and $P'(x) = [P(x)]^2 + xP(x)P'(x)$, we have $T_0(x) + xT_0'(x) = 2xP(x) + x^2P'(x)$. Note that $T_0(0) = 0$, solving this equation, we have $T_0(x) = xP(x)$. ■

In the following lemma, we give bijection proofs for $T_0(x) = xP(x)$ and $L(x) = x[P(x)]^2$.

Lemma 3.2. *For any $n \geq 1$, there is a bijection from the sets $\mathcal{P}_n^n$ (resp. $\mathcal{P}_n^1$) to $\mathcal{P}_{n-1}$ (resp. $\mathcal{P}_{n-1, n; \leq n}$). Furthermore, we have $T_0(x) = xP(x)$ and $L(x) = x[P(x)]^2$.*

Proof. For any $\alpha = (n, a_1, \dots, a_{n-1}) \in \mathcal{P}_n^n$, it is easy to check that $(a_1, \dots, a_{n-1}) \in \mathcal{P}_{n-1}$. Obviously, this is a bijection. Hence, $p_n^n = p_{n-1}$ for any $n \geq 1$. Similarly, for any

$\beta = (1, b_1, \dots, b_{n-1}) \in \mathcal{P}_n^1$, we have $(b_1, \dots, b_{n-1}) \in \mathcal{P}_{n-1, n; \leq n}$. Clearly, this is a bijection. So, $p_n^1 = p_{n-1, n; \leq n}$ for any $n \geq 1$.

By these two bijections, simple computations tell us that $T_0(x) = xP(x)$ and $L(x) = x[P(x)]^2$. ■

Theorem 3.3. *Let $k \geq 1$ and $T_k(x)$ be the generating function for the sequence p_n^{n-k} , then $T_k(x)$ satisfies the following recurrence relation*

$$T_k(x) = T_{k-1}(x) + \frac{(k+1)^{k-1}}{k!} x^{k+1} P(x) - \frac{2(k+1)^{k-2}}{(k-1)!} x^k,$$

equivalently,

$$T_k(x) = P(x) \sum_{i=0}^k \frac{(i+1)^{i-1}}{i!} x^{i+1} - \sum_{i=1}^k \frac{2(i+1)^{i-2}}{(i-1)!} x^i.$$

Proof. From Identity (7), we have

$$\sum_{n \geq k+1} \frac{p_n^{n-k}}{(n-1)!} x^n - \sum_{n \geq k+1} \frac{p_n^{n-k+1}}{(n-1)!} x^n = \frac{(k+1)^{k-1}}{k!} x^k \sum_{n \geq k+1} \frac{(n-k)^{n-k-2}}{(n-k-1)!} x^{n-k}$$

. Hence,

$$T_k(x) = T_{k-1}(x) + \frac{(k+1)^{k-1}}{k!} x^{k+1} P(x) - \frac{2(k+1)^{k-2}}{(k-1)!} x^k. ■$$

Corollary 3.4. *Let $T(x, y) = \sum_{k \geq 0} T_k(x) y^k$, then $T(x, y)$ satisfies the following equations*

$$\begin{aligned} (1-y) \left[T(x, y) + x \frac{\partial T(x, y)}{\partial y} \right] &= 2xP(x)P(xy) + x^2P(xy)P'(x) + x^2yP(x)P'(xy) \\ &\quad - 2xyP'(xy), \end{aligned}$$

and

$$T(x, y) - xP(x) = yT(x, y) + xP(x)[P(xy) - 1] - xy[P(xy)]^2.$$

The second identity implies that

$$T(x, y) = \frac{xP(xy)[P(x) - yP(xy)]}{1-y}.$$

Proof. By Theorem 3.3, we have

$$\sum_{k \geq 1} T_k(x)y^k = y \sum_{k \geq 0} T_k(x)y^k + xP(x) \sum_{k \geq 1} \frac{(k+1)^{k-1}}{k!} (xy)^k - \sum_{k \geq 1} \frac{2(k+1)^{k-2}}{(k-1)!} (xy)^k.$$

Since $T(x, y) = \sum_{k \geq 0} T_k(x)y^k$, we have $T(x, y) - T_0(x) = yT(x, y) + xP(x)[P(xy) - 1] - L(xy)$.

By Lemma 3.1, $T(x, y)$ satisfies the following equations

$$(1-y) \left[T(x, y) + x \frac{\partial T(x, y)}{\partial y} \right] = 2xP(x)P(xy) + x^2P(xy)P'(x) + x^2yP(x)P'(xy) - 2xyP'(xy),$$

and

$$T(x, y) - xP(x) = yT(x, y) + xP(x)[P(xy) - 1] - xy[P(xy)]^2.$$

Hence,

$$T(x, y) = \frac{xP(xy)[P(x) - yP(xy)]}{1-y}.$$

■

Given $s \geq k \geq 0$, we define a generating function $F_{k,s}(x) = \sum_{n \geq s+1} \frac{p_{n; \leq n-k}^{n-s}}{(n-1)!} x^n$, let $F_k(x, y) = \sum_{s \geq k} F_{k,s}(x)y^s$ and $F(x, y, z) = \sum_{k \geq 0} F_k(x, y)z^k$.

Theorem 3.4. *Let $s \geq k \geq 0$ and $F_{k,s}(x)$ and $T_s(x)$ be the generating function for $p_{n; \leq n-k}^{n-s}$ and p_n^{n-s} , respectively, then $F_{k,s}(x)$ satisfies the following recurrence relation*

$$F_{k,s}(x) = T_s(x) - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} x^i F_{k-i, s-i}(x) \quad (8)$$

for any $1 \leq k \leq s$, with the initial condition $F_{0,s}(x) = T_s(x)$. Furthermore, we have

$$F_{k,s}(x) = \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i T_{s-i}(x). \quad (9)$$

Proof. Taking $l = n - s$ in Identity (5) of Lemma 2.9, we have

$$p_{n; \leq n-k}^{n-s} = p_n^{n-s} - \sum_{i=1}^k \binom{n-1}{i} (k-i+1)(k+1)^{i-1} p_{n-i; \leq n-k}^{n-s}. \quad (10)$$

Note that $p_{n; \leq n}^{n-s} = p_n^{n-s}$. Hence, $F_{0,s}(x) = T_s(x) = \sum_{n \geq s+1} \frac{p_n^{n-s}}{(n-1)!} x^n$. By Identity (10), we have

$$\sum_{n \geq s+1} \frac{p_{n; \leq n-k}^{n-s}}{(n-1)!} x^n = \sum_{n \geq s+1} \frac{p_n^{n-s}}{(n-1)!} x^n - \sum_{n \geq s+1} \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} \frac{p_{n-i; \leq n-k}^{n-s}}{(n-i-1)!} x^n. \text{ Hence,}$$

$$F_{k,s}(x) = T_s(x) - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} x^i F_{k-i, s-i}(x)$$

We assume that $F_{k',s}(x) = \sum_{i=0}^{k'} \frac{(-1)^i (k'+1-i)^i}{i!} x^i T_{s-i}(x)$ for any $k' \leq k$, then

$$\begin{aligned}
F_{k+1,s}(x) &= T_s(x) - \sum_{i=1}^{k+1} \frac{(k-i+2)(k+2)^{i-1}}{i!} x^i \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} x^j T_{s-i-j}(x) \\
&= T_s(x) - \sum_{i=1}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i T_{s-i}(x) \sum_{j=1}^i \binom{i}{j} (k+2-j)(k+2)^{j-1} \left(-\frac{1}{k+2-m}\right)^j \\
&= \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} x^i T_{s-i}(x)
\end{aligned}$$

■

Corollary 3.5. *Let $k \geq 0$ and $F_k(x, y) = \sum_{s \geq k} F_{k,s}(x) y^s$, then $F_k(x, y)$ satisfies the following recurrence relation*

$$F_k(x, y) = T(x, y) - \sum_{i=0}^{k-1} T_i(x) y^i - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} (xy)^i F_{k-i}(x, y),$$

equivalently,

$$F_k(x, y) = T(x, y) \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i - \sum_{s=0}^{k-1} \sum_{i=0}^{k-1-s} \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i T_s(x) y^s.$$

Proof. Clearly, $F_0(x, y) = T(x, y)$. Identity (8) implies that

$$\sum_{s \geq k} F_{k,s}(x) y^s = \sum_{s \geq k} T_s(x) y^s - \sum_{s \geq k} \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} x^i F_{k-i,s-i}(x) y^s.$$

Hence,

$$F_k(x, y) = T(x, y) - \sum_{s=0}^{k-1} T_s(x) y^s - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} (xy)^i F_{k-i}(x, y),$$

Furthermore, by Identity (9), for any $k \geq 1$, we have

$$\begin{aligned}
F_k(x, y) &= \sum_{s \geq k} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} x^i T_{s-i}(x) y^s \\
&= \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i \left[T(x, y) - \sum_{s=0}^{k-1-i} T_s(x) y^s \right] \\
&= T(x, y) \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i - \sum_{s=0}^{k-1} \sum_{i=0}^{k-1-s} \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i T_s(x) y^s.
\end{aligned}$$

■

Corollary 3.6. Let $F(x, y, z) = \sum_{k \geq 0} F_k(x, y) z^k$, then

$$F(x, y, z) = \frac{T(x, y) - zT(x, yz)}{e^{xyz} - z}$$

Proof. By Corollary 3.5, we have

$$\begin{aligned} F(x, y, z) &= F_0(x, y) + \sum_{k \geq 1} F_k(x, y) z^k \\ &= T(x, y) \sum_{k \geq 0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i z^k \\ &\quad - \sum_{k \geq 1} \sum_{s=0}^{k-1} \sum_{i=0}^{k-1-s} \frac{(-1)^i (k+1-i)^i}{i!} (xy)^i T_s(x) y^s z^k \\ &= \frac{T(x, y) - zT(x, yz)}{e^{xyz} - z}. \end{aligned}$$

■

Given $s \geq k \geq 0$, we define a generating function $D_{k,s}(x) = \sum_{n \geq s+1} \frac{p_{n;n-k}^{n-s}}{(n-1)!} x^n$.

Corollary 3.7. Let $s \geq k \geq 0$ and $D_{k,s}(x)$ be the generating function for $p_{n;n-k}^{n-s}$. Then when $s = k$,

$$D_{k,k}(x) = \begin{cases} xP(x) & \text{if } k = 0 \\ x[P(x) - 1] & \text{if } k = 1 \\ P(x) \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=0}^{k-2} \frac{(-1)^i (k-1-i)^i}{i!} x^i & \text{if } k \geq 2 \end{cases}$$

When $s \geq k+1$, $D_{k,s}(x)$ satisfies the following recurrence relation

$$D_{k,s}(x) = D_{k-1,s}(x) + \frac{x^{k+1}}{(k+1)!} T_{s-k-1}(x) - \sum_{i=1}^k \frac{x^i}{i!} D_{k-i,s-i}(x),$$

with initial conditions $D_{0,0}(x) = xP(x)$ and $D_{0,s}(x) = T_{s-1}(x)$ for any $s \geq 1$. Furthermore, for any $k \geq 0$ and $s \geq k+1$, we have

$$D_{k,s}(x) = \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} T_{s-i}(x) [(k+1-i)^i - (k+2-i)^i].$$

Proof. Corollary 2.4 implies that $D_{0,0}(x) = xP(x)$ and $D_{k,k}(x) = x \left[R_{k-1}(x) - \frac{p_{k-1;\leq 0}}{(k-1)!} x^{k-1} \right]$ for any $k \geq 1$. Hence, $D_{1,1}(x) = x[P(x) - 1]$ since $p_{0;\leq 0} = 1$. When $k \geq 2$, $D_{k,k}(x) = xR_{k-1}(x)$ since $p_{k-1;\leq 0} = 0$. So, by Theorem 3.2, we have

$$D_{k,k}(x) = P(x) \sum_{i=0}^{k-1} \frac{(-1)^i (k-i)^i}{i!} x^i - \sum_{i=0}^{k-2} \frac{(-1)^i (k-1-i)^i}{i!} x^i$$

for any $k \geq 2$.

Furthermore, by Corollary 2.4, for any $k \geq 1$ and $s \geq k + 1$, we have

$$\sum_{n \geq s+1} \frac{p_{n;=n-k}^{n-s}}{(n-1)!} x^n = \sum_{n \geq s+1} \frac{p_{n;=n-k+1}^{n-s}}{(n-1)!} x^n + \sum_{n \geq s+1} \frac{\binom{n-1}{k+1} p_{n-k-1}^{n-s}}{(n-1)!} x^n - \sum_{n \geq s+1} \sum_{i=1}^k \frac{\binom{n-1}{i} p_{n-i;=n-k}^{n-s}}{(n-1)!} x^n.$$

Hence,

$$D_{k,s}(x) = D_{k-1,s}(x) + \frac{x^{k+1}}{(k+1)!} T_{s-k-1}(x) - \sum_{i=1}^k \frac{x^i}{i!} D_{k-i,s-i}(x).$$

For any $s \geq k \geq 1$, by Lemma 2.9, we have $p_{n;=n-k+1}^{n-s} = \sum_{i=1}^k \binom{n-1}{i} p_{n-i;\leq n-k}^{n-s}$, hence, $D_{k-1,s}(x) = \sum_{i=1}^k \frac{x^i}{i!} F_{k-i,s-i}(x)$. By Theorem 3.4, for any $k \geq 0$ and $s \geq k + 1$, we obtain

$$\begin{aligned} D_{k,s}(x) &= \sum_{i=1}^{k+1} \frac{x^i}{i!} F_{k+1-i,s-i}(x) \\ &= \sum_{i=1}^{k+1} \frac{x^i}{i!} \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} x^j T_{s-i-j}(x) \\ &= \sum_{i=1}^{k+1} \frac{x^i}{i!} T_{s-i}(x) \sum_{j=1}^i \binom{i}{j} (i-k-2)^{i-j} \\ &= \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} T_{s-i}(x) [(k+1-i)^i - (k+2-i)^i]. \end{aligned}$$

■

Corollary 3.8. *Let $k \geq 0$ and $D_k(x, y) = \sum_{s \geq k} D_{k,s}(x) y^s$, then $D_k(x, y)$ satisfies the following recurrence relation*

$$\begin{aligned} D_k(x, y) &= D_{k-1}(x, y) - D_{k-1,k-1}(x) y^{k-1} - D_{k-1,k}(x) y^k + D_{k,k}(x) y^k \\ &\quad + \frac{(xy)^{k+1}}{(k+1)!} T(x, y) - \sum_{i=1}^k \frac{(xy)^i}{i!} [D_{k-i}(x, y) - D_{k-i,k-i}(x) y^{k-i}] \end{aligned}$$

with initial condition $D_0(x) = xP(x) + T(x, y)$. Furthermore, we have

$$\begin{aligned} D_k(x) &= D_{k,k}(x) y^k + \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T(x, y) \\ &\quad - \sum_{s=0}^{k-1} \sum_{i=1}^{k-s} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T_s(x) y^s. \end{aligned}$$

Proof. By Corollary 3.7, it is easy to check that

$$D_0(x) = xP(x) + T(x, y).$$

Furthermore, for any $k \geq 1$, we have

$$\sum_{s \geq k+1} D_{k,s}(x)y^s = \sum_{s \geq k+1} D_{k-1,s}(x)y^s + \sum_{s \geq k+1} \frac{x^{k+1}}{(k+1)!} T_{s-k-1}(x)y^s - \sum_{s \geq k+1} \sum_{i=1}^k \frac{x^i}{i!} D_{k-i,s-i}(x)y^s.$$

Hence,

$$\begin{aligned} D_k(x, y) &= D_{k-1}(x, y) - D_{k-1,k-1}(x)y^{k-1} - D_{k-1,k}(x)y^k + D_{k,k}(x)y^k \\ &\quad + \frac{(xy)^{k+1}}{(k+1)!} T(x, y) - \sum_{i=1}^k \frac{(xy)^i}{i!} [D_{k-i}(x, y) - D_{k-i,k-i}(x)y^{k-i}]. \end{aligned}$$

On the other hand,

$$\begin{aligned} D_k(x, y) &= \sum_{s \geq k} D_{k,s}(x)y^s \\ &= D_{k,k}(x)y^k + \sum_{s \geq k+1} D_{k,s}(x)y^s \\ &= D_{k,k}(x)y^k + \sum_{s \geq k+1} \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} T_{s-i}(x) [(k+1-i)^i - (k+2-i)^i] y^s \\ &= D_{k,k}(x)y^k + \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] [T(x, y) - \sum_{s=0}^{k-i} T_s(x)y^s] \\ &= D_{k,k}(x)y^k + \sum_{i=1}^{k+1} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T(x, y) \\ &\quad - \sum_{s=0}^{k-1} \sum_{i=1}^{k-s} (-1)^i \frac{x^i}{i!} [(k+1-i)^i - (k+2-i)^i] T_s(x)y^s. \end{aligned}$$

■

Corollary 3.9. Let $D(x, y, z) = \sum_{k \geq 0} D_k(x, y)z^k$, then

$$D(x, y, z) = \frac{xe^{xyz}(P(x) - yz)}{e^{xyz} - yz} + \frac{e^{xyz} - 1}{z} F(x, y, z).$$

Proof. For any $s \geq k \geq 1$, by Lemma 2.9, we have $p_{n;=n-k+1}^{n-s} = \sum_{i=1}^k \binom{n-1}{i} p_{n-i; \leq n-k}^{n-s}$, hence,

$D_{k-1,s}(x) = \sum_{i=1}^k \frac{x^i}{i!} F_{k-i,s-i}(x)$. So,

$$\begin{aligned}
D_k(x, y) &= \sum_{s \geq k} D_{k,s}(x) y^s \\
&= D_{k,k}(x) y^k + \sum_{s \geq k+1} D_{k,s}(x) y^s \\
&= D_{k,k}(x) y^k + \sum_{s \geq k+1} \sum_{i=1}^{k+1} \frac{x^i}{i!} F_{k+1-i,s-i}(x) y^s \\
&= D_{k,k}(x) y^k + \sum_{i=1}^{k+1} \frac{(xy)^i}{i!} F_{k+1-i}(x, y).
\end{aligned}$$

Therefore,

$$\begin{aligned}
D(x, y, z) &= \sum_{k \geq 0} D_{k,k}(x) (yz)^k + \sum_{k \geq 0} \sum_{i=1}^{k+1} \frac{(xy)^i}{i!} F_{k+1-i}(x, y) z^k \\
&= \sum_{k \geq 0} D_{k,k}(x) (yz)^k + \sum_{k \geq 0} \sum_{i=0}^k \frac{(xy)^{i+1}}{(i+1)!} F_{k-i}(x, y) z^k \\
&= xP(x) - xyz + xyzR(x, yz) + \frac{e^{xyz} - 1}{z} F(x, y, z) \\
&= \frac{xe^{xyz}(P(x) - yz)}{e^{xyz} - yz} + \frac{e^{xyz} - 1}{z} F(x, y, z).
\end{aligned}$$

■

4 The asymptotic behavior of the sequences

In this section, we consider the asymptotic behavior of the sequences in previous sections.

First, we study the asymptotic behavior for $p_{n; \leq n-k}$. For a fixed k , let $\mu_k = \lim_{n \rightarrow \infty} \frac{p_{n; \leq n-k}}{(n+1)^{n-1}}$ and define a generating function $\mu(x) = \sum_{k \geq 0} \mu_k x^k$.

Theorem 4.1. *Let $k \geq 0$. The sequence μ_k satisfies the recurrence relation*

$$\mu_k = 1 - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} e^{-i} \mu_{k-i}$$

and $\mu_0 = 1$, equivalently,

$$\mu_k = \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i}.$$

Let $\mu(x)$ be the generating function for μ_k , then

$$\mu(x) = (e^{\frac{x}{e}} - x)^{-1}.$$

Proof. Clearly, $\mu_0 = 1$. Since we have used Identity (1) in Lemma 2.6 to prove Theorem 3.2, we want to obtain the results in Theorem 4.1 by Identity (2) in Lemma 2.6. For a fixed $i \geq 0$, we have $\lim_{n \rightarrow \infty} \binom{n}{i} \frac{(n-i+1)^{n-i-1}}{(n+1)^{n-1}} = \frac{e^{-i}}{i!}$. Hence,

$$\mu_k = 1 - \sum_{i=1}^k \frac{(k-i+1)(k+1)^{i-1}}{i!} e^{-i} \mu_{k-i}.$$

Now, we assume that $\mu_{k'} = \sum_{i=0}^{k'} \frac{(-1)^i (k'+1-i)^i}{i!} e^{-i}$ for any $k' \leq k$, then,

$$\begin{aligned} \mu_{k+1} &= 1 - \sum_{i=1}^{k+1} \frac{(k-i+2)(k+2)^{i-1}}{i!} e^{-i} \mu_{k+1-i} \\ &= 1 - \sum_{i=1}^{k+1} \frac{(k-i+2)(k+2)^{i-1}}{i!} e^{-i} \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} e^{-j} \\ &= 1 - \sum_{i=1}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i} \sum_{j=1}^i \binom{i}{j} (k+2-j)(k+2)^{j-1} \left(-\frac{1}{k+2-i}\right)^j \\ &= \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i}. \end{aligned}$$

Hence,

$$\begin{aligned} \mu(x) &= \sum_{k \geq 0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} x^k \\ &= \sum_{k \geq 0} \frac{(-1)^k e^{-k}}{k!} \sum_{i \geq 0} (i+1)^k x^{i+k} \\ &= (e^{\frac{x}{e}} - x)^{-1} \end{aligned}$$

■

For a fixed k , let $\eta_k = \lim_{n \rightarrow \infty} \frac{p_{n,i=n-k}}{(n+1)^{n-1}}$ and define a generating function $\eta(x) = \sum_{k \geq 0} \eta_k x^k$.

Corollary 4.1. $\eta_k = \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i}$ for any $k \geq 0$. Let $\eta(x)$ be the generating function for η_k , then $\eta(x) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)}$.

Proof. By Lemma 2.6, we have $\eta_{k-1} = \sum_{i=1}^k \frac{e^{-i}}{i!} \mu_{k-i}$ for any $k \geq 1$. Hence, by Theorem 4.1, for any $k \geq 0$,

$$\begin{aligned} \eta_k &= \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} e^{-j} \\ &= \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} - \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i}. \end{aligned}$$

Furthermore,

$$\eta(x) = \sum_{k \geq 0} \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \mu_{k+1-i} x^k = \frac{e^{\frac{x}{e}} - 1}{x} \mu(x) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)}$$

■

Now, we consider the asymptotic behavior for p_n^l . Given $l \geq 1$, let $\tau_l = \lim_{n \rightarrow \infty} \frac{p_n^l}{(n+1)^{n-2}}$. Clearly, $\tau_1 = 2$. We define a generating function $\tau(x) = \sum_{l \geq 1} \tau_l x^l$.

Lemma 4.1. *The sequence τ_l satisfies the recurrence relation*

$$\tau_l - \tau_{l+1} = \frac{l^{l-2}}{(l-1)!} e^{-l}$$

for any $l \geq 1$, with the initial condition $\tau_1 = 2$. Let $\tau(x)$ be the generating function for τ_l , then

$$\tau(x) = \frac{\frac{x^2}{e} P(\frac{x}{e}) - 2x}{x-1}.$$

Proof. For a fixed $l \geq 1$, we have $\lim_{n \rightarrow \infty} \frac{\binom{n-1}{l-1} (n-l+1)^{n-l-1}}{(n+1)^{n-2}} = \frac{e^{-l}}{(l-1)!}$. So, $\tau_l - \tau_{l+1} = \frac{l^{l-2}}{(l-1)!} e^{-l}$. Hence, $\sum_{l \geq 1} \tau_l x^l - \sum_{l \geq 1} \tau_{l+1} x^l = \sum_{l \geq 1} \frac{l^{l-2}}{(l-1)!} e^{-l} x^l$. $\tau(x)$ satisfies the equation $(x-1)\tau(x) + 2x = \frac{x^2}{e} P(\frac{x}{e})$. Equivalently, $\tau(x) = \frac{\frac{x^2}{e} P(\frac{x}{e}) - 2x}{x-1}$. ■

For fixed l and k , let $\rho_{l,k} = \lim_{n \rightarrow \infty} \frac{p_{n; \leq n-k}^l}{(n+1)^{n-2}}$ and define a generating function $\rho_l(x) = \sum_{k \geq 0} \rho_{l,k} x^k$.

Theorem 4.2. *Let $l \geq 1$, then the sequence $\rho_{l,k}$ satisfies the following recurrence relation*

$$\rho_{l,k} = \rho_{l,k-1} - \sum_{i=1}^k \frac{e^{-i}}{i!} \rho_{l,k-i}.$$

for any $k \geq 1$. Furthermore, we have $\rho_{l,k} = \rho_{l,0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i}$. Let $\rho_l(x)$ be the generating function for $\rho_{l,k}$, then $\rho_l(x) = \rho_{l,0} (e^{\frac{x}{e}} - x)^{-1}$. Furthermore, let $\rho(x, y) = \sum_{l \geq 1} \rho_l(x) y^l$, then

$$\rho(x, y) = \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{(y-1)(e^{\frac{x}{e}} - x)}$$

Proof. Since we have used Identity (5) in Lemma 2.9 to prove Theorem 3.4, we want to obtain the results in Theorem 4.2 by Identity (4) in Lemma 2.9. For a fixed $i \geq 0$, it is easy to obtained that $\lim_{n \rightarrow \infty} \binom{n-1}{i} \frac{(n-i+1)^{n-i-2}}{(n+1)^{n-2}} = \frac{e^{-i}}{i!}$. Hence,

$$\rho_{l,k} = \rho_{l,k-1} - \sum_{i=1}^k \frac{e^{-i}}{i!} \rho_{l,k-i}.$$

Now, assume that $\rho_{l,k'} = \rho_{l,0} \sum_{i=0}^{k'} \frac{(-1)^i (k'+1-i)^i}{i!} e^{-i}$ for any $k' \leq k$, then

$$\begin{aligned} \rho_{l,k+1} &= \rho_{l,k} - \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \rho_{l,k+1-i} \\ &= \rho_{l,0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} - \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \rho_{l,0} \sum_{j=0}^{k+1-i} \frac{(-1)^j (k+2-i-j)^j}{j!} e^{-j} \\ &= \rho_{l,0} \sum_{i=0}^{k+1} \frac{(-1)^i (k+2-i)^i}{i!} e^{-i} \end{aligned}$$

Hence, $\rho_l(x) = \sum_{k \geq 0} \rho_{l,0} \sum_{i=0}^k \frac{(-1)^i (k+1-i)^i}{i!} e^{-i} x^k = \rho_{l,0} (e^{\frac{x}{e}} - x)^{-1}$. Since $\rho_{l,0} = \pi_l$, by Lemma 4.1, we have

$$\begin{aligned} \rho(x, y) &= \sum_{l \geq 1} \pi_l (e^{\frac{x}{e}} - x)^{-1} y^l \\ &= \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{(y-1)(e^{\frac{x}{e}} - x)} \end{aligned}$$

■

For a fixed k , let $\lambda_{l,k} = \lim_{n \rightarrow \infty} \frac{p'_{n;n-k}}{(n+1)^{n-2}}$ and define a generating function $\lambda_l(x) = \sum_{k \geq 0} \lambda_{l,k} x^k$.

Corollary 4.2. *Let $\lambda_l(x)$ be the generating function for $\lambda_{l,k}$, then $\lambda_l(x) = \frac{e^{\frac{x}{e}} - 1}{x} \rho_l(x)$. Let $\lambda(x, y) = \sum_{l \geq 1} \lambda_l(x) y^l$, then $\lambda(x, y) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)} \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{y-1}$.*

Proof. Lemma 2.9 implies that $\lambda_{l,k} = \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \rho_{l,k+1-i}$. So, $\lambda_l(x) = \sum_{k \geq 0} \sum_{i=1}^{k+1} \frac{e^{-i}}{i!} \rho_{l,k+1-i} x^k = \frac{e^{\frac{x}{e}} - 1}{x} \rho_l(x)$. Hence, $\lambda(x, y) = \frac{e^{\frac{x}{e}} - 1}{x} \rho(x, y) = \frac{e^{\frac{x}{e}} - 1}{x(e^{\frac{x}{e}} - x)} \frac{\frac{y^2}{e} P(\frac{y}{e}) - 2y}{y-1}$.

5 Appendix

For convenience to check the identities in the previous sections, by the computer search, for $n \leq 7$, we obtain the number of parking functions $\mathcal{P}_{n;\leq s}^l$ and list them in Table 1. Note that $p_{n;\leq s}^l = 0$ if $l > s$.

	$l = 1$	2	3	4	5	6	7	8	$p_{n;\leq s}$
$(n, s) = (1, 1)$	1	0							1
$(2, 1)$	1	0							1
$(2, 2)$	2	1	0						3
$(3, 1)$	1	0							1
$(3, 2)$	4	3	0						7
$(3, 3)$	8	5	3	0					16
$(4, 1)$	1	0							1
$(4, 2)$	8	7	0						15
$(4, 3)$	26	19	16	0					61
$(4, 4)$	50	34	25	16	0				125
$(5, 1)$	1	0							1
$(5, 2)$	16	15	0						31
$(5, 3)$	80	65	61	0					206
$(5, 4)$	232	171	143	125	0				671
$(5, 5)$	432	307	243	189	125	0			1296
$(6, 1)$	1	0							1
$(6, 2)$	32	31	0						63
$(6, 3)$	242	211	206	0					659
$(6, 4)$	982	776	701	671	0				3130
$(6, 5)$	2642	1971	1666	1456	1296	0			9031
$(6, 6)$	4802	3506	2881	2401	1921	1296	0		16807
$(7, 1)$	1	0							1
$(7, 2)$	64	63	0						127
$(7, 3)$	728	665	659	0					2052
$(7, 4)$	4020	3361	3175	3130	0				13686
$(7, 5)$	14392	11262	10026	9351	9031	0			54062
$(7, 6)$	36724	27693	23667	20922	18682	16807	0		144495
$(7, 7)$	65536	48729	40953	35328	30208	24583	16807	0	262144

Table.1. The values of $p_{n;\leq s}^l$ for $1 \leq n \leq 7$

References

- [1] Sen-Peng Eu, Tung-Shan Fu, Chun-Ju Lai, On the enumeration of parking functions by leading terms, *Adv. in Appl. Math.* 35 (2005) 392-406
- [2] Sen-Peng Eu, Tung-Shan Fu, Yeong-Nan Yeh, Refined Chung-Feller theorems for lattice paths, *J. Combin. Theory, Ser. A* 112 (2005) 143-162.
- [3] Sen-Peng Eu, Shu-Chung Liu, Yeong-Nan Yeh, Taylor expansions for Catalan and Motzkin numbers, *Adv. in Appl. Math.* 29 (2002) 345-357.
- [4] J. Françon, Acyclic and parking functions, *J. Combin. Theory Ser. A* 18 (1975) 27-35.
- [5] D. Foata, J. Riordan, Mappings of acyclic and parking functions, *Aequationes Math.* 10 (1974) 10-22.
- [6] J.D. Gilbey, L.H. Kalikow, Parking functions, valet functions and priority queues, *Discrete Math.* 197/198 (1999) 351-373.
- [7] Po-Yi Huang, Jun Ma, Yeong-Nan Yeh, Ordered k -flaw Preference sets, submitted.
- [8] Po-Yi Huang, Jun Ma, Chun-Chen Yeh, k -flaw Preference sets, submitted.
- [9] J.P.S. Kung, C.H. Yan, Gončarov polynomials and parking functions, *J. Combin. Theory Ser. A* 102 (2003) 16C37.
- [10] J. Pitman, R. Stanley, A polytope related to empirical distributions, plane trees, parking functions, and the associahedron, *Discrete Comput. Geom.* 27 (4) (2002) 603-634.
- [11] J. Riordan, Ballots and trees, *J. Combin. Theory* 6 (1969) 408-411.
- [12] M.P. Schützenberger, On an enumeration problem, *J. Combin. Theory* 4 (1968) 219-221.
- [13] R.P. Stanley, Hyperplane arrangements, interval orders and trees, *Proc. Natl. Acad. Sci.* 93 (1996) 2620-2625.
- [14] R.P. Stanley, Parking functions and non-crossing partitions, in: The Wilf Festschrift, *Electron. J. Combin.* 4 (1997) R20.

- [15] C.H. Yan, Generalized tree inversions and k-parking functions, *J. Combin. Theory Ser. A* 79 (1997) 268-280.
- [16] C.H. Yan, On the enumeration of generalized parking functions, *Congr. Numer.* 147 (2000) 201-209.
- [17] C.H. Yan, Generalized parking functions, tree inversions and multicolored graphs, *Adv. in Appl. Math.* 27 (2001) 641-670.